arXiv:1103.2744v1 [cond-mat.str-el] 14 Mar 2011

Ward Identities for the Two-Coupled-Chains Model

L. C. Costa

Institut Für Theoretische Physik, Universität Frankfurt
Max-von-Laue Strasse 1, 60438 Frankfurt am Main, Germany.
Centro de Ciências Naturais e Humanas,
Universidade Federal do ABC, 09210-170, Santo André, Brazil.

A. Ferraz

International Institute of Physics, Universidade Federal
do Rio Grande do Norte, 59012-970, Natal, Brazil.

V. Mastropietro

Dipartamento di Matematica, Università di Roma "Tor Vergata"
V. le Ricerca Scientifica, 00133, Roma, Italy.

Abstract

We derive exact Ward Identities in the Two-Coupled-Chains Model in the presence of a momentum cutoff, with an extra correction term generated by the breaking of local symmetries. Such term is non vanishing in 1-loop order in the infinite bandwidth limit and this indicates the presence of a quantum anomaly.

PACS numbers: 05.30.Fk, 05.30.Rt

I. INTRODUCTION

The two-coupled chains model (TCCM) is a quasi-one-dimensional system of interacting fermions located along two Tomonaga-Luttinger chains coupled by a interchain hopping (usually denoted by $t_\perp$). In the last two decades the TCCM has also been investigated as a prototype model to test the possibility of having a higher dimensional Luttinger liquid state [1]-[3]. Moreover there were important predictions indicating that the TCCM might also manifest quantum confinement regime [4]-[6] at strong coupling. This quantum transition occurs when the electronic motion becomes strictly one dimensional and, as a result of the interaction, the renormalized interchain hopping $t_\perp^R$ becomes nullified. Since the strong coupling regime is very difficult to deal with even in such a simplified system, this problem remains open to this date, even if several perturbative and non-perturbative methods have been used to treat the TCCM [7]-[11].

A very powerful method in the analysis of low dimensional fermionic systems is bosonization, used by Mattis and Lieb [12] in their exact solution of the Tomonaga-Luttinger chain, a model obtained from a system of non relativistic fermions linearizing the energy relations in the vicinity of the Fermi points (see Tomonaga [13]). The presence of the Dirac sea generates *anomalous density commutators* of the associated Kac-Moody algebra in the Tomonaga-Luttinger model which are at the core of the bosonization method. Indeed, it was well known, as was pointed out earlier by Jordan [14] and Schwinger [15], that when the number of degrees of freedom in the ground state of a physical system becomes infinite, anomalous non-zero commutators of currents or density operators are naturally produced by such a state.

However, the solvability of the Tomonaga-Luttinger chain is due to certain special features of the system which allow us to reduce it into harmonic oscillator modes. This feature is absent in more realistic models, like the TCCM model, at least for general interactions. One way to proceed is to combine approximate bosonization schemes with the Renormalization Group method (see [16]). Another possible promising approach is based on the combination of Renormalization Group methods with the Ward Identities (WI), which must however be derived with care. Indeed even in the single chain problem the presence of anomalous commutators reflect themselves in the presence of quantum anomalies in the WI; the (formal) WI derived neglecting the ultraviolet (UV) cutoffs which are necessary to regularize the theory are not correct and extra terms must be added to it. This was pointed out by Johnson [17] and Georgi and Rawls [18] (in order to verify the Adler result in d=1+1 [19]), by deriving the correct WI for the massless Thirring model making use of the exact solution. The presence of anomalies in the WI on the other hand can be easily overlooked in a formal diagrammatic approach, as in the Dzyaloshinkii and Larkin solution of the Tomonaga-Luttinger model [20] (see also [7]); an anomaly is present in the infinite bandwidth limit and the non-regularized WI is no longer valid in that case.

The derivation of the WI for the TCCM is more tricky, as in such a case there is no exact solution from which the WI can be derived, and the presence of anomalies can be easily overlooked. In this paper we derive, in section II, the (non-regularized) WI associated with the existing chiral symmetry of the TCCM proceeding formally from the functional integrals with no UV cutoff; in section III we show that such an identity is *violated*, by an explicit one-loop perturbative calculation. We then consider in section IV the theory with an ultraviolet momentum cutoff and we derive, following [21], the WI with an extra term generated by the presence of the cutoff which actually breaks the local invariance of the model. This extra term in the WI is written as a functional integral and in Section V we show that its lowest order contribution in the WI is *non vanishing* even when the UV cutoff is taken to infinity; this signals the presence of an anomaly in the TCMM in this limit.

Benfatto and Mastropietro obtained for systems in the Tomonaga-Luttinger model universality class in [21] (see also [22],[23]) a full non-perturbative control of this extra term in the WI in the limit of a removed UV cutoff, using the methods of Constructive Quantum Field Theory; it turns out that such a correction is proportional, in the limit of a removed cutoff, to the vertex function and, in the case of the Tomonaga-Luttinger model, it reproduces the same WI given by the anomalous commutators.

In the case of the TCCM only if the backscattering and the umklapp interactions are neglected the model becomes solvable and the WI correction term reduces to the Tomonaga-Luttinger model result; with general interactions there is no reason for this to happen. While the non-perturbative limit of the correcting term in the Tomonaga-Lutinger model is probably technically quite hard to implement, we believe that this correction term, which is found here for the first time, plays an important role in the physical properties of the TCCM.

II. THE TWO-COUPLED-CHAINS MODEL

The TCCM describes two spinless Tomonaga-Luttinger chains coupled together by interchain hopping. After diagonalization the spinless electrons are distributed among the bonding (b) and anti-bonding (a) bands with a Fermi surface (FS) consisting of four points, as displayed in our Figure 1. The corresponding bare free action S_f for single particles located in the vicinities of those Fermi points of the TCCM is, in this way, given by

$$S_f = \sum_{\substack{j=a,b \\ \alpha=\pm}} \int_{\mathrm{x}} {\psi^{+}}^{j}_{\alpha}(\mathrm{x})(i(\partial_t + \alpha v_F^j \partial_x) + \gamma^j)\psi^j_\alpha(\mathrm{x}), \tag{1}$$

where $\mathrm{x} = (t, x)$, the flavor index j refers to the bonding and antibonding bands and the index α denotes the right (+) and left (-) sides of the FS. Here v_F^j is the Fermi velocity, k_F^j is the Fermi wave vector and $\gamma^j = v_F^j k_F^j$ is a characteristic energy for j-th "flavor", used to clarify Figure 1. Following common practice among workers in the field [4] we next define the corresponding bare interacting action S_{int} by



FIG. 1: Fermi surface representation of TCCM, formed by the location of four bare momenta $\pm k_F^a$ (antibonding band) and $\pm k_F^b$ (boundig band).

$$\begin{aligned} S_{int} \;=\; & \sum_{j,\alpha} \int_{\mathrm{x},\mathrm{x}'} \Bigg\{ g_0 {\psi^{+}}^{j}_{\alpha}(\mathrm{x}) {\psi^{+}}^{j}_{-\alpha}(\mathrm{x}') \psi^j_{-\alpha}(\mathrm{x}') \psi^j_\alpha(\mathrm{x}) + \sum_{i\neq j} \Bigg[g_{bs} {\psi^{+}}^{j}_{\alpha}(\mathrm{x}) {\psi^{+}}^{i}_{-\alpha}(\mathrm{x}') \psi^j_{-\alpha}(\mathrm{x}') \psi^i_\alpha(\mathrm{x}) + \\ & + \; g_f {\psi^{+}}^{j}_{\alpha}(\mathrm{x}) {\psi^{+}}^{i}_{-\alpha}(\mathrm{x}') \psi^i_{-\alpha}(\mathrm{x}') \psi^j_\alpha(\mathrm{x}) + g_u {\psi^{+}}^{j}_{\alpha}(\mathrm{x}) {\psi^{+}}^{j}_{-\alpha}(\mathrm{x}') \psi^i_{-\alpha}(\mathrm{x}') \psi^i_\alpha(\mathrm{x}) \Bigg] \Bigg\}, \end{aligned} \tag{2}$$

with $g_0 \equiv g_0(x-x')$ being the intraband forward short range interaction, and $g_{bs} \equiv g_{bs}(x-x')$ (backscattering), $g_f \equiv g_f(x-x')$ (forward) and $g_u \equiv g_u(x-x')$ (umklapp) the three interband short range interactions. See Figure 2, for illustration. Notice that although g_u (g_{bs}) doesn't preserve flavor globally (locally), S_{int} is fully symmetric under chiral (+,−) transformation. If g_u and g_{bs} are vanishing the model becomes solvable and equivalent to a Tomonaga-

Luttinger model, while in the presence of non vanishing g_u and g_{bs} an exact solution is unknown.



FIG. 2: Diagrammatic representation for the TCCM interactions for right- and left-electrons, respectively, with $j \neq j'$. Full and dashed lines represent $+$ and $-$ electrons, respectively, in the vicinities of k_F^j and $-k_F^j$ respectively.

Using the total bare action $S = S_f + S_{int}$, we define next the functional generators Z and W by

$$Z[\xi, \xi^+, J] = \mathrm{e}^{iW[\xi,\xi^+,J]} = \int [d\psi^+][d\psi] \ \exp\left[i(S_f + S_{int} + \xi \cdot \psi + \psi^+ \cdot \xi^+ + J \cdot \psi^+\psi)\right], \quad (3)$$

where, for simplicity, we use the notation $\xi \cdot \psi = \sum_{j,\alpha} \int_{\mathrm{x}} \ \xi_\alpha^j(\mathrm{x})\psi_\alpha^j(\mathrm{x})$.

Let us next apply the local phase transformation to the fermionic fields:

$$\psi_\alpha^j(\mathrm{x}) \to e^{-i\lambda(\mathrm{x})}\psi_\alpha^j(\mathrm{x}), \quad (4)$$

$$\psi^{+j}_{\ \alpha}(\mathrm{x}) \to e^{i\lambda(\mathrm{x})}\psi^{+j}_{\ \alpha}(\mathrm{x}). \quad (5)$$

Under these $U(1)$ transformations, the action S becomes

$$S \to S - \sum_{\alpha,j} \int_{\mathrm{x}} \lambda(\mathrm{x})(\partial_t + \alpha v_F^j \partial_x)\psi^{+j}_{\ \alpha}(\mathrm{x})\psi_\alpha^j(\mathrm{x}), \quad (6)$$

and the source terms transform as

$$\xi \cdot \psi \to \xi \cdot \psi - i \sum_{\alpha,j} \int_{\mathrm{x}} \lambda(\mathrm{x})\xi_\alpha^j(\mathrm{x})\psi_\alpha^j(\mathrm{x}), \quad (7)$$

$$\psi^+ \cdot \xi^+ \to \psi^+ \cdot \xi^+ + i \sum_{\alpha,j} \int_{\mathrm{x}} \lambda(\mathrm{x})\psi^{+j}_{\ \alpha}(\mathrm{x})\xi^{+j}_{\ \alpha}(\mathrm{x}). \quad (8)$$

Since these phase transformations leave Z invariant, we immediately arrive at the WIs

$$\sum_{j,\alpha} \left[i(\partial_t + \alpha v_F^j \partial_x)\frac{\delta}{\delta J_\alpha^j(\mathrm{x})} - \xi_\alpha^j(\mathrm{x})\frac{\delta}{\delta \xi_\alpha^j(\mathrm{x})} + \xi^{+j}_{\ \alpha}(\mathrm{x})\frac{\delta}{\delta \xi^{+j}_{\ \alpha}(\mathrm{x})} \right] Z = 0, \quad (9)$$

or, equivalently

$$\sum_{j,\alpha}\left[i(\partial_t+\alpha v_F^j\partial_x)\frac{\delta}{\delta J_\alpha^j(\mathrm{x})}-\xi_\alpha^j(\mathrm{x})\frac{\delta}{\delta\xi_\alpha^j(\mathrm{x})}+\bar{\xi}_\alpha^j(\mathrm{x})\frac{\delta}{\delta\bar{\xi}_\alpha^j(\mathrm{x})}\right]W=0. \tag{10}$$

Alternatively we can rewrite the WIs in terms of $\Gamma[\langle\psi^+\rangle,\langle\psi\rangle,J]$, the generator of the one-particle irreducible functions. Defining

$$\Gamma[\langle\psi\rangle,\langle\psi^+\rangle,J]=W[J,\xi,\bar{\xi}]-{\xi^+}_\alpha^j\cdot\langle\psi_\alpha^j\rangle+\langle{\psi^+}_\alpha^j\rangle\cdot\xi_\alpha^j, \tag{11}$$

with

$$\langle\psi_\alpha^j(\mathrm{x})\rangle=\frac{\delta W}{\delta{\xi^+}_\alpha^j(\mathrm{x})},\qquad\qquad \langle{\psi^+}_\alpha^j(\mathrm{x})\rangle=-\frac{\delta W}{\delta\xi_\alpha^j(\mathrm{x})}, \tag{12}$$

we find

$$\sum_{j,\alpha}i(\partial_t+\alpha v_F^j\partial_x)\frac{\delta\Gamma}{\delta J_\alpha^j(\mathrm{x})}=\sum_{j,\alpha}\left(\langle\bar{\psi}_\alpha^j(\mathrm{x})\rangle\frac{\delta}{\delta\langle{\psi^+}_\alpha^j(\mathrm{x})\rangle}-\langle\psi_\alpha^j(\mathrm{x})\rangle\frac{\delta}{\delta\langle\psi_\alpha^j(\mathrm{x})\rangle}\right)\Gamma. \tag{13}$$

Consequently, if we differenciate this last equation with respect to

$$\frac{\delta^2}{\delta\langle\psi_\beta^k(\mathrm{x}')\rangle\delta\langle{\psi^+}_\beta^k(\mathrm{x}'')\rangle}, \tag{14}$$

we obtain, after setting the fields equal to zero,

$$\begin{aligned}\sum_{j,\alpha}\ i\ \ (\partial_t+\alpha v_F^j\partial_x)&\frac{\delta^3\Gamma}{\delta\langle\psi_\beta^k(\mathrm{x}')\rangle\delta\langle{\psi^+}_\beta^k(\mathrm{x}'')\rangle\delta J_\alpha^j(\mathrm{x})}=\\ &=\ \delta(\mathrm{x}-\mathrm{x}')\frac{\delta^2\Gamma}{\delta\langle\psi_\beta^k(\mathrm{x})\rangle\delta\langle{\psi^+}_\beta^k(\mathrm{x}'')\rangle}-\delta(\mathrm{x}-\mathrm{x}'')\frac{\delta^2\Gamma}{\delta\langle\psi_\beta^k(\mathrm{x}')\rangle\delta\langle{\psi^+}_\beta^k(\mathrm{x})\rangle},\end{aligned} \tag{15}$$

which, in momenta space, reduces to the more conventional form

$$\sum_{\alpha,j}\left(q_0-\alpha v_F^jq\right)\Gamma_{\beta,\alpha}^{k,j(2,1)}(\mathrm{p}+\mathrm{q},\mathrm{p})=\Gamma_\beta^{k(2)}(\mathrm{p}+\mathrm{q})-\Gamma_\beta^{k(2)}(\mathrm{p}), \tag{16}$$

with $\mathrm{p}=(p_0,p)$ and $\Gamma_\beta^{k(2)}(\mathrm{p})=(G_\beta^k(\mathrm{p}))^{-1}$ being the inverse of the (k,β)-th single particle propagator. Writing

$$\Gamma_{\beta,\alpha}^{k,j(2,1)}(\mathrm{p}+\mathrm{q},\mathrm{p})=\delta_j^k\delta_\alpha^\beta+\Lambda_{\beta,\alpha}^{k,j}(\mathrm{p}+\mathrm{q},\mathrm{p}), \tag{17}$$

this can be expressed in a more convenient form:

$$\sum_{\alpha,j}(q_0-\alpha v_F^jq)\Lambda_{\beta,\alpha}^{k,j}(\mathrm{p}+\mathrm{q},\mathrm{p})=\Sigma_\beta^k(\mathrm{p})-\Sigma_\beta^k(\mathrm{p}+\mathrm{q}), \tag{18}$$

where $\Sigma_\beta^k(\mathrm{p})$ is the self-energy associated with the (k,β)-th particle.

III. PERTURBATIVE TEST OF THE WARD IDENTITY

The above derivation of the WI is purely formal; the functional integrals have just a formal meaning so that the results of their manipulations may not be true. This is precisely what happens here since, as we can see from perturbation theory, this WI is no longer valid. In zero-th order the previously derived WI is trivially satisfyed. Choosing, for convenience, $k = b$ and $\beta = +$, we have simply in the zero-th order:

$$\Gamma_{+}^{b(2)}(\mathrm{p}+\mathrm{q}) - \Gamma_{+}^{b(2)}(\mathrm{p}) = q_0 - v_F^b q \tag{19}$$

and

$$\Gamma_{+,\alpha}^{b,j}(\mathrm{p}+\mathrm{q},\mathrm{q}) = \delta_{+,\alpha}\delta_{b,j}, \tag{20}$$

in agreement with equation (16). In 1-loop order there are just two tadpole contribuctions to the self-energy, as indicated in Figure (3).



FIG. 3: One-loop self-energy diagrams for a bonding electron in the k_F^b branch. The couplings are g_0 and g_f, respectively.

It is straightforward to calculate these two diagrams and we immediatly find

$$\Sigma_{+}^{b}(\mathrm{p}) - \Sigma_{+}^{b}(\mathrm{p}+\mathrm{q}) = 0. \tag{21}$$

There are two vertex function in 1-loop order associated with those self-energy diagrams: $\Lambda_{+,-}^{b,b}$ and $\Lambda_{+,-}^{b,a}$. They are displayed in Figure (4).

Calculating their contributions we find

$$\Lambda_{+,-}^{b,b}(\mathrm{p}+\mathrm{q},\mathrm{p}) = \frac{g_0(q)}{2\pi}|q|G_{-}^{b}(\mathrm{q}), \tag{22}$$

$$\Lambda_{+,-}^{b,a}(\mathrm{p}+\mathrm{q},\mathrm{p}) = \frac{g_f(q)}{2\pi}|q|G_{-}^{a}(\mathrm{q}), \tag{23}$$

with G_{-}^{b} (G_{-}^{a}) being the noninteracting Green's function for the bonding (anti-bonding) spinless "$-$" electrons. As a result it follows from that

$$(q_0 + v_F^b q)\Lambda_{+,-}^{b,b} + (q_0 + v_F^a q)\Lambda_{+,-}^{b,a} = \frac{|q|}{2\pi}(g_0(q) + g_f(q)), \tag{24}$$



FIG. 4: The two vertices corresponding to the self-energies diagrams of FIG.3.

in clear disagreement with our WI, for $q \neq 0$. If we take the ultraviolet limit $\Lambda \to \infty$ this WI violation reflects the chiral symmetry breaking produced by the Dirac seas for the bonding and anti-bonding bands. However if Λ is finite our derivation of the WI should also be modified to take the presence of such a cuttoff explicitly into account. We will do that next.

IV. WARD IDENTITY IN THE PRESENCE OF THE CUTOFF

The WI in the presence of a finite cutoff for the Luttinger liquid was obtained by Benfatto and Mastropietro [21] and here we adopt their derivation for our purposes, considering imaginary times for convenience. Let us therefore define the functional generator W in the presence of the cutoff Λ as

$$\exp(-W) = \int \mathcal{D}[\psi^+\psi]\text{exp } \left(S_\Lambda + \xi^+ \cdot \psi + \psi^+ \cdot \xi + J \cdot \psi^+\psi\right). \tag{25}$$

The cutoff Λ manifests itself in the regulators $C^\alpha_j(\mathrm{x},\Lambda)$'s which now appears in our definition of the non-interacting action S^Λ_f:

$$S^\Lambda_f = \sum_{\alpha,j} \int_{\mathrm{k}} \psi^{+j}_{\alpha}(\mathrm{k})(C^j_\alpha(k,\Lambda)^{-1}(ik_0 - \alpha v^j_F k + \gamma^j)\psi^j_\alpha(\mathrm{k})), \tag{26}$$

where $C^j_\alpha(p;\Lambda)$ can either be a step function or any other smoother function we wish. This non-interacting action is now consistent with the definition of a non-interacting single particle propagator $G^j_\alpha(\mathrm{p})$

$$G^j_\alpha(\mathrm{p}) = \frac{C^j_\alpha(p;\Lambda)}{-ip_0 + \alpha v_F(p - \alpha k^j_F)}.$$

If we now follow the derivation of the WI as before and work in energy-momentum space we find

$$\int \mathcal{D}[\psi^+,\psi]\sum_{\alpha,j}\left\{(iq_0-\alpha v_F^j q)\int\limits_{\mathrm{k}}\psi^{+j}_{\alpha}(\mathrm{k}+\mathrm{q})\psi^j_\alpha(\mathrm{k})+\int\limits_{\mathrm{k}}\xi^{+j}_{\alpha}(\mathrm{k}+\mathrm{q})\psi^j_\alpha-\int\limits_{\mathrm{k}}\psi^{+j}_{\alpha}(\mathrm{k}+\mathrm{q})\xi^j_\alpha(\mathrm{k})\right.$$
$$\left.+\int\limits_{\mathrm{k}}\psi^{+j}_{\alpha}(\mathrm{k}+\mathrm{q})(C^j_\alpha(\mathrm{k},\mathrm{q}))^{-1}\psi^j_\alpha(\mathrm{k})\right\}\exp i(S+\xi^+\cdot\psi+\psi\cdot\xi+J\cdot\psi\psi)=0, \tag{27}$$

where $\int\limits_{\mathrm{k}}=(2\pi)^{-2}\int dk_0\int dk$ and

$$\begin{aligned}(C^j_\alpha(\mathrm{k},\mathrm{q}))^{-1} \;=\;& \left[(C^j_\alpha(k+q,\Lambda))^{-1}-1\right]\left[i(k_0+q_0)-\alpha v_F^j(k+q)\right]-\\ -\;& \left[(C^j_\alpha(k,\Lambda))^{-1}-1\right]\left[ik_0-\alpha v_F^j k\right].\end{aligned} \tag{28}$$

That is, the presence of the cutoff produces a new Λ-dependent term in the WI. For convenience we introduce a new source term to our functional generator W_Λ, namely

$$\sum_{\alpha,j}\int\limits_{\mathrm{q},\mathrm{k}}\chi^j_\alpha(q)\psi^{+j}_{\alpha}(\mathrm{k}+\mathrm{q})(C^j_\alpha(k,q))^{-1}\psi^j_\alpha(\mathrm{k}). \tag{29}$$

In this way, our generalized cutoff dependent WI becomes

$$\begin{aligned}&\sum_{\alpha,j}\;(iq_0-\alpha v_F^j q)\;\frac{\delta W_\Lambda}{\delta J^j_\alpha(\mathrm{q})}\\ &\qquad=\qquad\sum_{\alpha,j}\left\{\int_{\mathrm{k}}\left(\xi^j_\alpha(\mathrm{k})\frac{\delta}{\delta\xi^{\mathrm{j}}_\alpha(\mathrm{k}+\mathrm{q})}-\xi^{+\mathrm{j}}_{\alpha}(\mathrm{k})\frac{\delta}{\delta\xi^{+\mathrm{j}}_{\alpha}(\mathrm{k}+\mathrm{q})}\right)-\frac{\delta}{\delta\chi^j_\alpha(\mathrm{q})}\right\}W_\Lambda.\end{aligned} \tag{30}$$

Defining the 1PI functional generator Γ as before we can, equivalently, rewrite this WI as

$$\begin{aligned}&\sum_{\alpha,j}\qquad(iq_0-\alpha v_F^j q)\frac{\delta\Gamma}{\delta J^j_\alpha(\mathrm{q})}\\ &\quad=\sum_{\alpha,j}\left\{\int_{\mathrm{k}}\left(\langle\psi^j_\alpha(\mathrm{k})\rangle\frac{\delta}{\delta\langle\psi^{\mathrm{j}}_\alpha(\mathrm{k}+\mathrm{q})\rangle}-\langle\psi^{+\mathrm{j}}_{\alpha}(\mathrm{k})\rangle\frac{\delta}{\delta\langle\psi^{+\mathrm{j}}_{\alpha}(\mathrm{k}+\mathrm{q})\rangle}\right)-\frac{\delta}{\delta\chi^j_\alpha(\mathrm{q})}\right\}\Gamma.\end{aligned} \tag{31}$$

If we functionally differentiate this last result with respect

$$\frac{\delta^2}{\delta\langle\psi^j_\alpha(\mathrm{k}')\rangle\delta\langle\psi^{+j}_{\alpha}(\mathrm{k}'')\rangle}, \tag{32}$$

we obtain

$$\sum_{\alpha',j'}\left(iq_0-\alpha' v_F^{j'}q\right)\Gamma^{j,j'(2,1)}_{\alpha,\alpha'}(\mathrm{k}+\mathrm{q},\mathrm{k})=\Gamma^{j(2)}_\alpha(\mathrm{k}+\mathrm{q})-\Gamma^{j(2)}_\alpha(\mathrm{k})+\sum_{\alpha',j'}\Delta^{j,j'}_{\alpha,\alpha'}(\mathrm{k},\mathrm{q}), \tag{33}$$

where

$$\Delta^{j,j'}_{\alpha,\alpha'}(\mathrm{k},\mathrm{q})=-\frac{\delta^3\Gamma}{\delta\langle\psi^j_\alpha(\mathrm{k})\rangle\delta\langle\psi^{+j}_{\alpha}(\mathrm{k}+\mathrm{q})\rangle\delta\chi^{j'}_{\alpha'}(\mathrm{q})}. \tag{34}$$

These Δ's are precisely the new terms produced by the finite cutoff Λ.

V. WARD IDENTITY RESTAURATION IN THE PRESENCE OF A CUTOFF

In the previous section we derived the exact WI for the theory with a cutoff; the functional integrals are well defined and the throughout resulting WI differs with respect to its non-regularized counterpart by a correcting term. Of course, now the WI should be true order by order in perturbation theory. Indeed let us calculate this cutoff contribution in one-loop order. Since at this order we have that



FIG. 5: $\Delta^{j,j'}_{+,-}$ in one-loop order.

$$(C^{j'}_{-}(\mathrm{k}',\mathrm{q},\Lambda))^{-1} = -(iq_0+v^{j'}_F q)+(C^{j'}_{-}(k'+q))^{-1}(i(k'_0+q_0)+v^{j'}_F(k'+q))-(C^{j'}_{-}(k))^{-1}(ik'_0+v^{j'}_F k'), \tag{35}$$

it follows immediately that

$$\sum_{j'}\int_{k'} \quad \left[(C^{j'}_{-}(\mathrm{k}'+\mathrm{q}))^{-1}\left[i(k'_0+q_0)+v^{j'}_F(k'+q)\right]-(C^{j'}_{-}(k))^{-1}\left[ik'_0+v^{j'}_F k'\right]\right] G^{j'}_{-}(\mathrm{k}')G^{j'}_{-}(\mathrm{k}'+\mathrm{q})$$
$$= \sum_{j'}\int_{k'}\left[G^{j'}_{-}(\mathrm{k}')-G^{j'}_{-}(\mathrm{k}'+\mathrm{q})\right]=0. \tag{36}$$

As a result, we obtain the trivial identity

$$\sum_{j'=b,a}(iq_0+v^{j'}_F q)\Lambda^{b,j'}_{+,-} = \sum_{j'=b,a}(iq_0+v^{j'}_F q)\Lambda^{b,j'}_{+,-} \tag{37}$$

and, as expected, the WI is fully restored for a finite cutoff. In the limit $\Lambda \to \infty$, $\Lambda^{b,j'}_{+,-}$ is given by (22),(23) times an extra i. Note that the lowest order contribution to $\Delta^{jj'}_{\pm}$ is non vanishing even in the limit $\Lambda \to \infty$. A genuine anomaly is present in this case since the regularized WI differs from its non-regularized counterpart. Note also that, even if the lowest order contribution of $\Delta^{jj'}_{\pm}$ depends only on g_0 and g_f, the higher order terms depend on the other couplings as well.

VI. CONCLUSION

In the absence of backscattering and umklapp interactions the TCCM can be solved exactly by bosonization methods but such interactions cannot be mapped into a quadratic bosonic expression; therefore no solution exists in the general case. In order to derive the WI, we perform first a phase transformation in the functional integrals with no cutoff to derive identities connecting the vertex function with the self energy. The non-regularized formal WI's obtained in this way, even if widely accepted, are however not always true, as an explicit one-loop calculation demonstrates. Indeed, without regularizations the functional integrals are ill-defined and one should not take for granted the WI which result from their manipulation. On the other hand, if a momentum cutoff is imposed from the start the functional integrals are well defined and an exact WI is derived, with a new extra term $\Delta_{\pm}^{jj'}$ with respect to the non-regularized result. This extra term can be written as a functional integral and a perturbative computation shows that its first order contribution is already *non* vanishing, signaling the presence of the anomaly in the infinite bandwidth limit.

Note that the correction term $\Delta_{\pm}^{jj'}$ differs from result obtained for the Tomonaga-Luttinger model, since it depends on *all* the interactions appearing in the TCCM. Only if g_b and g_u are nullified one can apply the non perturbative analysis of [21] which gives $\Delta_{\pm}^{jj'}$ proportional to the vertex function, in the limit of a removed cutoff, in agreement with the result obtained from the anomalous commutators. In the general case, however, there is no reason to expect that the WI for the TCCM has the same form as in the Tomonaga-Luttinger model. While the non-perturbative limit of the correcting term in the Tomonaga-Luttinger model is probably technically quite hard to analyze, we believe that this correcting term, plays an important role in the physical properties of the TCCM as well.

Acknowledgments

We acknowledge the comments we had from E. Kochetov and P. Kopietz. LCC would also thanks G. Bonneau, L. Bartosch, A. Isidori, A. Kreisel and E. Correa for many valuable discussion. This work was supported by the CNPq, CAPES (Probral) and by the Brazilian

Min. of Science and Technology.

[1] P. W. Anderson, *Science* **235**, 1196 (1987).

[2] P. W. Anderson, Phys. Rev. Lett. **67**, 3844 (1991).

[3] D. G. Clarke, S Strong and P. W. Anderson, Phys. Rev. Lett. **74**, 4499 (1995).

[4] M. Fabrizio, Phys. Rev. B **48**, 15838 (1993).

[5] S. Ledowski, P. Kopietz and A. Ferraz, Phys. Rev. B **71**, 235103 (2005).

[6] S. Ledowski and P. Kopietz, Phys. Rev. B **75**, 045134 (2007).

[7] J. Sólyom, Adv. in Phys. **28**, 201 (1979).

[8] C. Bourbonnais and L. G. Caron, Int. J. Mod. Phys. B **5**, 1033 (1991).

[9] C. Castellani, C. Di Castro and W. Metzner, Phys. Rev. Lett. **69**, 1703 (1992).

[10] A. M. Finkel'stein and A. I. Larkin **47**, 10461 (1993).

[11] M. Fabrizio and A. Parola, Phys. Rev. Lett. **70**, 226 (1993).

[12] D. C. Mattis and E. H. Lieb, J. Math. Phys. **6**, 304 (1965).

[13] S. Tomonaga, Progr. Theoret. Phys. **92**, 609 (1950).

[14] P. Jordan, Z. Physik **93**, 464 (1935); **98**, 759 (1936); **99**, 109 (1936); **102**, 243 (1936); **105**, 114 (1937); **105** 229 (1937).

[15] J. Schwinger, Phys. Rev. Lett. **3**, 296 (1959).

[16] J. Voit, Phys. Rev. B **45**, 4027 (1992).

[17] K. Johnson, Nuovo Cimento **20**, 4, 773 (1961).

[18] H. Georgi and J. M. Rawls, Phys. Rev. D **3**, 874 (1971).

[19] S. L. Adler and W. Bardeen, Phys. Rev. **182**, 1459 (1969).

[20] I. E. Dzyaloshinskii and A. I. Larkin, Sov. Phys JETP, **38**, 1, 202 (1973).

[21] G. Benfatto and V. Mastropietro, Comm. Math. Phys. **258**, 609 (2005).

[22] V. Mastropietro, J. Math. Phys. **48**, 022302, (2007).

[23] G. Benfatto V. Mastropietro and P. Falco , Comm. Math. Phys. **273**, 67-118 (2007).